

## Lesson 4.4 Exercises, pages 314–321

### A

3. For each function below, determine possible functions  $f$  and  $g$  so that  $y = f(g(x))$ .

a)  $y = (x + 4)^2$

**Sample solution:**

Let  $f(g(x)) = (x + 4)^2$

Replace  $x + 4$  with  $x$ .

Then,  $g(x) = x + 4$  and  $f(x) = x^2$

b)  $y = \sqrt{x + 5}$

**Sample solution:**

Let  $f(g(x)) = \sqrt{x + 5}$

Replace  $x + 5$  with  $x$ .

Then,  $g(x) = x + 5$  and  $f(x) = \sqrt{x}$

c)  $y = \frac{1}{x - 2}$

**Sample solution:**

Let  $f(g(x)) = \frac{1}{x - 2}$

Replace  $x - 2$  with  $x$ .

Then,  $g(x) = x - 2$  and  $f(x) = \frac{1}{x}$

d)  $y = (6 - x)^3$

**Sample solution:**

Let  $f(g(x)) = (6 - x)^3$


Replace  $6 - x$  with  $x$ .

Then,  $g(x) = 6 - x$  and  $f(x) = x^3$


**B**

4. Given  $f(x) = x + 3$  and  $g(x) = x^2 + 1$ , sketch the graph of each composite function below then state its domain and range.

a)  $y = f(f(x))$



b)  $y = f(g(x))$




Make a table of values for the functions.


$x$	$f(x)$	$f(f(x))$	$g(x)$	$f(g(x))$	$g(f(x))$	$g(g(x))$
-4	-1	2	17	20	2	290
-3	0	3	10	13	1	101
-2	1	4	5	8	2	26
-1	2	5	2	5	5	5
0	3	6	1	4	10	2
1	4	7	2	5	17	5
2	5	8	5	8	26	26

- a) Graph the points with coordinates  $(x, f(f(x)))$  that fit on the grid. Draw a line through the points for the graph of  $y = f(f(x))$ . From the graph, the domain is  $x \in \mathbb{R}$  and the range is  $y \in \mathbb{R}$ .
- b) Graph the points with coordinates  $(x, f(g(x)))$  that fit on the grid. Draw a smooth curve through the points for the graph of  $y = f(g(x))$ . From the graph, the domain is  $x \in \mathbb{R}$  and the range is  $y \geq 4$ .

c)  $y = g(f(x))$



d)  $y = g(g(x))$



- c) Graph the points with coordinates  $(x, g(f(x)))$  that fit on the grid. Draw a smooth curve through the points for the graph of  $y = g(f(x))$ . From the graph, the domain is  $x \in \mathbb{R}$ . From the table, the range is  $y \geq 1$ .
- d) Graph the points with coordinates  $(x, g(g(x)))$  that fit on the grid. Draw a smooth curve through the points for the graph of  $y = g(g(x))$ . From the graph, the domain is  $x \in \mathbb{R}$ . From the table, the range is  $y \geq 2$ .

5. Consider the function  $h(x) = (x - 1)(x + 5)$ .

- a) Why is it incorrect to write  $h(x) = f(g(x))$ , where  $f(x) = x - 1$  and  $g(x) = x + 5$ ?

**It is incorrect because, as written,  $h(x)$  is the product of  $f(x)$  and  $g(x)$ , not their composition.**

- b) For what functions  $f(x)$  and  $g(x)$  is  $h(x)$  a composite function?

**Expand:  $h(x) = (x - 1)(x + 5)$**

$$h(x) = x^2 + 4x - 5$$

**Complete the square:  $h(x) = (x^2 + 4x + 4) - 9$**

$$h(x) = (x + 2)^2 - 9$$

**Possible functions are:  $f(x) = x^2 - 9$  and  $g(x) = x + 2$  for  $h(x) = f(g(x))$**

6. For each pair of functions below:

- Determine an explicit equation for the indicated composite function.
- State the domain of the composite function, and explain any restrictions on the variable.

- a)  $f(x) = \sqrt{x + 1}$  and  $g(x) = x^2 - x - 6$ ;  $g(f(x))$

- i) In  $g(x) = x^2 - x - 6$ , replace  $x$  with  $\sqrt{x + 1}$ .

$$g(f(x)) = (\sqrt{x + 1})^2 - \sqrt{x + 1} - 6$$

$$g(f(x)) = x + 1 - \sqrt{x + 1} - 6$$

$$g(f(x)) = x - 5 - \sqrt{x + 1}$$

- ii) The domain of  $f(x) = \sqrt{x + 1}$  is  $x \geq -1$ .

The domain of  $g(x) = x^2 - x - 6$  is  $x \in \mathbb{R}$ .

So, the domain of  $g(f(x))$  is  $x \geq -1$ .

The variable  $x$  is restricted because the square root of a real number is only defined for numbers that are greater than or equal to 0.

- b)  $f(x) = \sqrt{x - 1}$  and  $g(x) = \frac{1}{x + 3}$ ;  $g(f(x))$

- i) In  $g(x) = \frac{1}{x + 3}$ , replace  $x$  with  $\sqrt{x - 1}$ .

$$g(f(x)) = \frac{1}{\sqrt{x - 1} + 3}$$

- ii) The domain of  $f(x) = \sqrt{x - 1}$  is  $x \geq 1$ .

The domain of  $g(x) = \frac{1}{x + 3}$  is  $x \neq -3$ .

$-3$  is not in the range of  $f(x)$ .

So, the domain of  $g(f(x))$  is  $x \geq 1$ .

The variable  $x$  is restricted because the square root of a real number is only defined for numbers that are greater than or equal to 0.

c)  $f(x) = \sqrt{x + 3}$  and  $g(x) = 2x - 1$ ;  $f(g(x))$

i) In  $f(x) = \sqrt{x + 3}$ , replace  $x$  with  $2x - 1$ .

$$f(g(x)) = \sqrt{2x - 1 + 3}$$

$$f(g(x)) = \sqrt{2x + 2}$$

ii) The domain of  $g(x) = 2x - 1$  is  $x \in \mathbb{R}$ .

The domain of  $f(x) = \sqrt{x + 3}$  is  $x \geq -3$ .

$$\text{So, } g(x) \geq -3$$

$$2x - 1 \geq -3$$

$$2x \geq -2$$

$$x \geq -1$$

So, the domain of  $f(g(x))$  is  $x \geq -1$ .

The variable  $x$  is restricted because the square root of a real number is only defined for numbers that are greater than or equal to 0.

d)  $f(x) = \frac{1}{x - 1}$  and  $g(x) = x^2 + 2x$ ;  $f(f(x))$

i) In  $f(x) = \frac{1}{x - 1}$ , replace  $x$  with  $\frac{1}{x - 1}$ .

$$f(f(x)) = \frac{1}{\frac{1}{x - 1} - 1}, \text{ which simplifies to } f(f(x)) = \frac{x - 1}{2 - x}, x \neq 1$$

ii) The domain of  $f(x) = \frac{1}{x - 1}$  is  $x \neq 1$ .

$$\text{Also, } 2 - x \neq 0$$

$$x \neq 2$$

So, the domain of  $f(f(x))$  is  $x \neq 1$  and  $x \neq 2$ .

The variable  $x$  is restricted because the denominator of a fraction can never be 0.

7. For each function below

i) Determine possible functions  $f$  and  $g$  so that  $y = f(g(x))$ .

ii) Determine possible functions  $f$ ,  $g$ , and  $h$  so that  $y = f(g(h(x)))$ .

a)  $y = x^2 - 6x + 5$

b)  $y = -3x^2 - 30x - 40$

Sample solution:

$$y = x^2 - 6x + 5$$

$$y = (x^2 - 6x + 9) - 4$$

$$y = (x - 3)^2 - 4$$

$$\text{Let } f(g(x)) = (x - 3)^2 - 4$$

i) Replace  $x - 3$  with  $x$ .

$$\text{Then, } g(x) = x - 3 \text{ and}$$

$$f(x) = x^2 - 4$$

ii) Replace  $x - 3$  with  $x$ .

$$\text{Then, } h(x) = x - 3, g(x) = x^2,$$

$$\text{and } f(x) = x - 4$$

Sample solution:

$$y = -3x^2 - 30x - 40$$

$$y = -3(x^2 + 10x + 25) + 75 - 40$$

$$y = -3(x + 5)^2 + 35$$

$$\text{Let } f(g(x)) = -3(x + 5)^2 + 35$$

i) Replace  $x + 5$  with  $x$ .

$$\text{Then, } g(x) = x + 5 \text{ and}$$

$$f(x) = -3x^2 + 35$$

ii) Replace  $x + 5$  with  $x$ .

$$\text{Then, } h(x) = x + 5, g(x) = x^2,$$

$$\text{and } f(x) = -3x + 35$$

$$\text{c) } y = \sqrt{(x - 2)^2 + 3}$$

Sample solution:

$$\text{Let } f(g(x)) = \sqrt{(x - 2)^2 + 3}$$

i) Replace  $x - 2$  with  $x$ .

Then,  $g(x) = x - 2$  and

$$f(x) = \sqrt{x^2 + 3}$$

ii) Replace  $x - 2$  with  $x$ .

Then,  $h(x) = x - 2$ ,

$$g(x) = x^2, \text{ and}$$

$$f(x) = \sqrt{x + 3}$$

$$\text{d) } y = \sqrt{x^2 + 4x + 3}$$

Sample solution:

$$y = \sqrt{x^2 + 4x + 3}$$

$$y = \sqrt{(x^2 + 4x + 4) - 1}$$

$$y = \sqrt{(x + 2)^2 - 1}$$

$$\text{Let } f(g(x)) = \sqrt{(x + 2)^2 - 1}$$

i) Replace  $x + 2$  with  $x$ .

Then,  $g(x) = x + 2$  and

$$f(x) = \sqrt{x^2 - 1}$$

ii) Replace  $x + 2$  with  $x$ .

Then,  $h(x) = x + 2$ ,  $g(x) = x^2$ ,

$$\text{and } f(x) = \sqrt{x - 1}$$

8. Create composite functions using either or both functions in each pair of functions below. In each case, how many different composite functions could you create? Justify your answer.

a)  $f(x) = |x|$  and  $g(x) = \frac{1}{x}$

$$f(f(x)) = ||x||, \text{ which simplifies to } f(f(x)) = |x|$$

$$f(g(x)) = \left| \frac{1}{x} \right|, \text{ which simplifies to } f(g(x)) = \frac{1}{|x|}$$

$$g(f(x)) = \frac{1}{|x|}$$

$$g(g(x)) = \frac{1}{\frac{1}{x}}, \text{ which simplifies to } g(g(x)) = x, x \neq 0$$

There are only 3 different composite functions, because  $f(g(x)) = g(f(x))$ .

b)  $f(x) = \sqrt{x}$  and  $g(x) = |x|$

$$f(f(x)) = \sqrt{\sqrt{x}}$$

$$f(g(x)) = \sqrt{|x|}$$

$$g(f(x)) = |\sqrt{x}|, \text{ which simplifies to } g(f(x)) = \sqrt{x}$$

$$g(g(x)) = ||x||, \text{ which simplifies to } g(g(x)) = |x|$$

There are 4 different composite functions.

c)  $f(x) = x^3$  and  $g(x) = \frac{1}{x}$

$$f(f(x)) = (x^3)^3, \text{ which simplifies to } f(f(x)) = x^9$$

$$f(g(x)) = \left( \frac{1}{x} \right)^3, \text{ which simplifies to } f(g(x)) = \frac{1}{x^3}$$

$$g(f(x)) = \frac{1}{x^3}$$

$$g(g(x)) = \frac{1}{\frac{1}{x}}, \text{ which simplifies to } g(g(x)) = x, x \neq 0$$

There are only 3 different composite functions, because  $f(g(x)) = g(f(x))$ .

9. Given the function  $y = \frac{x}{\sqrt{x-3}}$ , determine possible functions:

a)  $f$  and  $g$  so that  $y = \frac{f(x)}{g(x)}$

**Sample solution:**

$$f(x) = x \text{ and } g(x) = \sqrt{x-3}$$

b)  $f$ ,  $g$ , and  $h$  so that  $y = \frac{f(x)}{g(h(x))}$

**Sample solution:**

Replace  $x - 3$  with  $x$ .

Let  $h(x) = x - 3$ , then  $g(x) = \sqrt{x}$ , and  $f(x) = x$ .

c)  $f$  and  $g$  so that  $y = f(g(x))$

**Sample solution:**

When  $g(x)$  replaces  $x$  in  $f(x)$ , the numerator must be  $x$  and the denominator

must be  $\sqrt{x-3}$ . So,  $g(x) = x - 3$  and  $f(x) = \frac{x+3}{\sqrt{x}}$

10. Given the functions  $f(x) = \sqrt{x}$ ,  $g(x) = x^2 - x + 6$ , and  $k(x) = \frac{2}{x}$ , write an explicit equation for each combination.

a)  $h(x) = f(g(x)) + k(x)$

For  $f(g(x))$ , replace  $x$  in

$$f(x) = \sqrt{x} \text{ with } x^2 - x + 6.$$

$$\text{Then, } f(g(x)) = \sqrt{x^2 - x + 6}$$

$$\text{So, } h(x) = \sqrt{x^2 - x + 6} + \frac{2}{x},$$

$$x \neq 0$$

b)  $h(x) = g(f(x)) - f(g(x))$

For  $g(f(x))$ , replace  $x$  in

$$g(x) = x^2 - x + 6 \text{ with } \sqrt{x}.$$

$$\text{Then, } g(f(x)) = (\sqrt{x})^2 - \sqrt{x} + 6$$

$$\text{Or, } g(f(x)) = x - \sqrt{x} + 6, x \geq 0$$

$$\text{So, } h(x) = x - \sqrt{x} + 6 - \sqrt{x^2 - x + 6}, x \geq 0$$

c)  $h(x) = k(g(x)) + k(f(x))$

For  $k(g(x))$ , replace  $x$  in

$$k(x) = \frac{2}{x} \text{ with } x^2 - x + 6.$$

$$\text{Then, } k(g(x)) = \frac{2}{x^2 - x + 6}$$

For  $k(f(x))$ , replace  $x$  in

$$k(x) = \frac{2}{x} \text{ with } f(x) = \sqrt{x}$$

$$\text{Then, } k(f(x)) = \frac{2}{\sqrt{x}}, x > 0$$

$$\text{So, } h(x) = \frac{2}{x^2 - x + 6} + \frac{2}{\sqrt{x}}, x > 0$$

d)  $h(x) = f(g(x)) \cdot k(x)$

From part a,

$$f(g(x)) = \sqrt{x^2 - x + 6}$$

$$\text{So, } h(x) = \sqrt{x^2 - x + 6} \cdot \left(\frac{2}{x}\right), x \neq 0$$

**11.** Given the function  $y = (x^2 - 9)\sqrt{x + 2}$ , determine possible functions in each case:

a) functions  $f$  and  $g$  so that  $y = f(x) \cdot g(x)$

**Sample solution:**

$$f(x) = x^2 - 9 \text{ and } g(x) = \sqrt{x + 2}$$

b) functions  $f$ ,  $g$ , and  $h$  so that  $y = f(x) \cdot g(h(x))$

**Sample solution:**

$$f(x) = x^2 - 9$$

$$\text{For } g(h(x)), \text{ let } h(x) = x + 2, \text{ then } g(x) = \sqrt{x}$$

c) functions  $f$ ,  $g$ ,  $h$ , and  $k$  so that  $y = f(x) \cdot k(x) \cdot g(h(x))$

**Sample solution:**

$$\text{From part b, for } g(h(x)), \text{ let } h(x) = x + 2, \text{ then } g(x) = \sqrt{x}$$

$$\text{Factor: } x^2 - 9 = (x + 3)(x - 3)$$

$$\text{Then, } f(x) = x + 3 \text{ and } k(x) = x - 3$$

**12.** Is there a function  $f(x)$  such that each relationship is true?

Justify your answer.

a)  $f(f(x)) = f(x)$

b)  $f(f(x)) = f(x) + f(x)$

**Yes, when  $f(x) = x$ , then  $f(f(x)) = x$**

**Yes, when  $f(x) = 2x$ , then  $f(f(x)) = 4x$  and  $f(x) + f(x) = 2x + 2x$ , or  $4x$**

### C

**13.** Given  $f(x) = \frac{1}{x-2}$ ,  $g(x)$  is a quadratic function, and  $h(x) = f(g(x))$ , determine an explicit equation for  $g(x)$  for each situation below.

Explain your strategies.

a) The domain of  $h(x)$  is  $x \in \mathbb{R}$ .

**Sample solution: The denominator of  $h(x)$  must never be 0.**

$$\text{When } g(x) = x^2 + 3, \text{ then } f(g(x)) = \frac{1}{x^2 + 3 - 2}, \text{ which simplifies to}$$

$$f(g(x)) = \frac{1}{x^2 + 1}.$$

b) The domain of  $h(x)$  is  $x \neq a$  and  $x \neq b$ , where  $a$  and  $b$  are real numbers.

**Sample solution: There must be exactly two values of  $x$  that make the denominator of  $h(x)$  equal to 0. When  $g(x) = x^2 + 1$ , then**

$$f(g(x)) = \frac{1}{x^2 + 1 - 2}, \text{ which simplifies to } f(g(x)) = \frac{1}{x^2 - 1}.$$

$$\text{So, } a = 1 \text{ and } b = -1$$

c) The domain of  $h(x)$  is  $x \neq c$ , where  $c$  is a real number.

**Sample solution:** There must be exactly one value of  $x$  that makes the denominator of  $h(x)$  equal to 0. When  $g(x) = x^2 + 2$ , then

$$f(g(x)) = \frac{1}{x^2 + 2 - 2}, \text{ which simplifies to } f(g(x)) = \frac{1}{x^2}. \text{ So, } c = 0$$

**14.** Use  $f(x) = \frac{1-x}{1+x}$ .

a) Determine an explicit equation for  $f(f(x))$ , then state the domain of the function.

In  $f(x) = \frac{1-x}{1+x}$ , replace  $x$  with  $\frac{1-x}{1+x}$

$$\begin{aligned} f(f(x)) &= \frac{1 - \frac{1-x}{1+x}}{1 + \frac{1-x}{1+x}} \\ &= \frac{\frac{1+x - (1-x)}{1+x}}{\frac{1+x + (1-x)}{1+x}} \\ &= x, x \neq -1 \end{aligned}$$

The domain of the function is:  $x \neq -1$

b) What is the inverse of  $f(x)$ ? Explain.

Since  $f(f(x)) = x, x \neq -1$ , then  $f(x)$  is its own inverse.

So, the inverse of  $f(x)$  is  $f^{-1}(x) = \frac{1-x}{1+x}$ .