

PRACTICE TEST, pages 162–164

1. **Multiple Choice** Which statement is false?

A. $|x| = \sqrt{x^2}$ for $x \in \mathbb{R}$

B. $x = \sqrt{x^2}$ for $x \geq 0$

C. $|x| = \sqrt[3]{x^3}$ for $x \in \mathbb{R}$

D. $x = \sqrt[3]{x^3}$ for $x \geq 0$

2. **Multiple Choice** Which is the correct simplification of $\sqrt{12x^3}$?

A. $2\sqrt{3x^3}$, $x \geq 0$

B. $2x\sqrt{3x}$, $x \in \mathbb{R}$

C. $2x\sqrt{3x}$, $x \geq 0$

D. $2|x|\sqrt{3x}$, $x \in \mathbb{R}$

3. Simplify.

a) $\sqrt{72} - 5\sqrt{2} + 3\sqrt{8}$

$$\begin{aligned} &= \sqrt{36 \cdot 2} - 5\sqrt{2} + 3\sqrt{4 \cdot 2} \\ &= 6\sqrt{2} - 5\sqrt{2} + 6\sqrt{2} \\ &= 7\sqrt{2} \end{aligned}$$

b) $(\sqrt{3} - \sqrt{5})(\sqrt{3} + \sqrt{5})$

$$\begin{aligned} &= \sqrt{3}(\sqrt{3} + \sqrt{5}) - \sqrt{5}(\sqrt{3} + \sqrt{5}) \\ &= 3 + \sqrt{15} - \sqrt{15} - 5 \\ &= -2 \end{aligned}$$

c) $(2\sqrt{6} + 3\sqrt{5})^2$

$$\begin{aligned} &= (2\sqrt{6} + 3\sqrt{5})(2\sqrt{6} + 3\sqrt{5}) \\ &= 2\sqrt{6}(2\sqrt{6} + 3\sqrt{5}) \\ &\quad + 3\sqrt{5}(2\sqrt{6} + 3\sqrt{5}) \\ &= 24 + 6\sqrt{30} + 6\sqrt{30} + 45 \\ &= 69 + 12\sqrt{30} \end{aligned}$$

d) $\frac{1}{\sqrt{7} - \sqrt{3}}$

$$\begin{aligned} &= \frac{1}{\sqrt{7} - \sqrt{3}} \cdot \frac{(\sqrt{7} + \sqrt{3})}{(\sqrt{7} + \sqrt{3})} \\ &= \frac{\sqrt{7} + \sqrt{3}}{(\sqrt{7})^2 - (\sqrt{3})^2} \\ &= \frac{\sqrt{7} + \sqrt{3}}{4} \end{aligned}$$

4. Identify the values of the variable for which each radical is defined where necessary, then simplify.

a) $(\sqrt{x} + 2)(\sqrt{x} - 3)$

The radicands cannot be negative, so $x \geq 0$.

$$\begin{aligned} &(\sqrt{x} + 2)(\sqrt{x} - 3) \\ &= \sqrt{x}(\sqrt{x} - 3) + 2(\sqrt{x} - 3) \\ &= x - 3\sqrt{x} + 2\sqrt{x} - 6 \\ &= x - \sqrt{x} - 6 \end{aligned}$$

b) $6\sqrt{a^2} + 2a$

$\sqrt{a^2} \in \mathbb{R}$ when $a^2 \geq 0$.
 $a^2 \geq 0$, so $\sqrt{a^2}$ is defined for $a \in \mathbb{R}$.

$$\begin{aligned} &6\sqrt{a^2} + 2a \\ &= 6|a| + 2a \end{aligned}$$

$$\text{c) } \frac{8\sqrt{2}}{\sqrt{12} - \sqrt{10}}$$

$$= \frac{8\sqrt{2}}{2\sqrt{3} - \sqrt{10}}$$

$$= \frac{8\sqrt{2}}{(2\sqrt{3} - \sqrt{10})} \cdot \frac{(2\sqrt{3} + \sqrt{10})}{(2\sqrt{3} + \sqrt{10})}$$

$$= \frac{8\sqrt{2}(2\sqrt{3}) + 8\sqrt{2}(\sqrt{10})}{(2\sqrt{3})^2 - (\sqrt{10})^2}$$

$$= \frac{16\sqrt{6} + 8\sqrt{20}}{12 - 10}$$

$$= \frac{16\sqrt{6} + 16\sqrt{5}}{2}$$

$$= 8\sqrt{6} + 8\sqrt{5}$$

$$\text{d) } \frac{2\sqrt{10} - \sqrt{3}}{\sqrt{10} + \sqrt{3}}$$

$$= \frac{(2\sqrt{10} - \sqrt{3}) \cdot (\sqrt{10} - \sqrt{3})}{(\sqrt{10} + \sqrt{3}) \cdot (\sqrt{10} - \sqrt{3})}$$

$$= \frac{2\sqrt{10}(\sqrt{10} - \sqrt{3}) - \sqrt{3}(\sqrt{10} - \sqrt{3})}{(\sqrt{10})^2 - (\sqrt{3})^2}$$

$$= \frac{20 - 2\sqrt{30} - \sqrt{30} + 3}{10 - 3}$$

$$= \frac{23 - 3\sqrt{30}}{7}$$

5. Which equations have real roots? If the root is real, determine its value. If the equation has no real roots, explain how you know.

$$\text{a) } \sqrt{2x + 3} = 3$$

$$2x + 3 \geq 0; \text{ that is, } x \geq -\frac{3}{2}$$

$$\sqrt{2x + 3} = 3$$

$$(\sqrt{2x + 3})^2 = 3^2$$

$$2x + 3 = 9$$

$$2x = 6$$

$$x = 3$$

$x = 3$ lies in the set of possible values for x . So, the equation has a real root.

$$\text{b) } \sqrt{5x - 1} = \sqrt{2x + 5}$$

$$5x - 1 \geq 0; \text{ that is, } x \geq \frac{1}{5}$$

$$2x + 5 \geq 0; \text{ that is, } x \geq -\frac{5}{2}$$

So, for both radicals to be defined, $x \geq \frac{1}{5}$

$$\sqrt{5x - 1} = \sqrt{2x + 5}$$

$$(\sqrt{5x - 1})^2 = (\sqrt{2x + 5})^2$$

$$5x - 1 = 2x + 5$$

$$3x = 6$$

$$x = 2$$

$x = 2$ lies in the set of possible values for x . So, the equation has a real root.

$$\text{c) } \sqrt{3x + 2} + 5 = 2$$

$$3x + 2 \geq 0; \text{ that is, } x \geq -\frac{2}{3}$$

$$\sqrt{3x + 2} + 5 = 2$$

$$\sqrt{3x + 2} = -3$$

The left side of the equation is greater than or equal to 0.

The right side of the equation is negative, -3 .

So, no real solutions are possible.

The equation has no real roots.

$$\text{d) } 2\sqrt{x - 8} = 3\sqrt{x + 2}$$

$$x - 8 \geq 0; \text{ that is, } x \geq 8$$

$$x + 2 \geq 0; \text{ that is, } x \geq -2$$

So, for both radicals to be defined, $x \geq 8$

$$2\sqrt{x - 8} = 3\sqrt{x + 2}$$

$$(2\sqrt{x - 8})^2 = (3\sqrt{x + 2})^2$$

$$4(x - 8) = 9(x + 2)$$


$$4x - 32 = 9x + 18$$

$$-50 = 5x$$

$$x = -10$$

$x = -10$ does not lie in the set of possible values for x . So, the equation has no real roots.

6. To make a picture frame, a square with area 40 cm^2 is cut from a square with area 90 cm^2 . Serena wants to put a thin gold ribbon around the inside and outside edges of the frame. How much ribbon does Serena need?



The side length of a square is the square root of its area.

So, the side length of the square with area 40 cm^2 is:

$$\sqrt{40} = 2\sqrt{10} \text{ cm}$$

The side length of the square with area 90 cm^2 is:

$$\sqrt{90} = 3\sqrt{10} \text{ cm}$$

The length of ribbon needed = perimeter of large square
+ perimeter of small square

$$= 4(2\sqrt{10}) + 4(3\sqrt{10})$$

$$= 8\sqrt{10} + 12\sqrt{10}$$

$$= 20\sqrt{10}$$

Serena needs $20\sqrt{10} \text{ cm}$ of ribbon.

7. The formula $t = \sqrt{\frac{2d}{9.8}}$ gives the time, t seconds, for an object at rest to fall d metres. It took 2.5 s for a ball dropped from a roof to hit the ground. To the nearest metre, from what height was the ball dropped?

$$t = \sqrt{\frac{2d}{9.8}} \quad \text{Substitute: } t = 2.5$$

$$2.5 = \sqrt{\frac{2d}{9.8}}$$

$$(2.5)^2 = \left(\sqrt{\frac{2d}{9.8}}\right)^2$$

$$6.25 = \frac{2d}{9.8}$$

$$61.25 = 2d$$

$$d = 30.625$$

The ball was dropped from a height of about 31 m.